



LEARNING TO

INJECT *HEMLIBRA*

What is HEMLIBRA?

HEMLIBRA is a prescription medicine used for routine prophylaxis to prevent or reduce the frequency of bleeding episodes in adults and children, ages newborn and older, with hemophilia A with or without factor VIII inhibitors.

What is the most important information I should know about HEMLIBRA?

HEMLIBRA increases the potential for your blood to clot. People who use activated prothrombin complex concentrate (aPCC; Feiba®) to treat breakthrough bleeds while taking HEMLIBRA may be at risk of serious side effects related to blood clots.

These serious side effects include:

- **Thrombotic microangiopathy (TMA)**, a condition involving blood clots and injury to small blood vessels that may cause harm to your kidneys, brain, and other organs

- **Blood clots (thrombotic events)**, which may form in blood vessels in your arm, leg, lung, or head

Carefully follow your healthcare provider's instructions regarding when to use an on-demand bypassing agent or factor VIII, and the dose and schedule to use for breakthrough bleed treatment. If aPCC (Feiba®) is needed, talk to your healthcare provider in case you feel you need more than 100 U/kg of aPCC (Feiba®) total.

Your body may make antibodies against HEMLIBRA, which may stop HEMLIBRA from working properly. Contact your healthcare provider immediately if you notice that HEMLIBRA has stopped working for you (eg, increase in bleeds).

Please see Important Safety Information, including **Serious Side Effects**, throughout this brochure, as well as the accompanying HEMLIBRA full Prescribing Information and Medication Guide.



HEMLIBRA[®]
emicizumab-kxwh | 150
injection for subcutaneous use | mg/mL



As you begin your treatment with HEMLIBRA, this brochure will show you how to prepare and inject your medicine properly. Whether you're injecting yourself or a person you care for, this information is a reminder of what you learned at your healthcare provider's office or hemophilia treatment center. If you have any questions, be sure to ask your healthcare provider.

INSIDE, YOU'LL FIND INFORMATION ON:

Getting ready

Important information before you get started..... 5

Supplies you'll need 6

Preparing for your injection 8

Transferring HEMLIBRA from vial to syringe..... 10

Injecting HEMLIBRA 14

Combining vials..... 17

Disposing of your supplies 21

Important tips to remember..... 22

Please remember:

You should not inject yourself, or anyone else, unless you have been shown how to by your healthcare provider. If you have any questions, always ask your healthcare provider, and be sure you read and understand the Instructions for Use provided with your HEMLIBRA before you inject.

The information contained in this brochure does not provide medical advice or take the place of speaking with your healthcare provider about your medical condition or treatment.

What are the other possible side effects of HEMLIBRA?

The most common side effects of HEMLIBRA include: injection site reactions (redness, tenderness, warmth, or itching at the site of injection), headache, and joint pain. These are not all of the possible side effects of HEMLIBRA. You can speak with your healthcare provider for more information.

What else should I know about HEMLIBRA?

See the detailed "Instructions for Use" that comes with your HEMLIBRA for information on how to prepare and inject a dose of HEMLIBRA, and how to properly throw away (dispose of) used needles and syringes.

- Stop taking your prophylactic bypassing therapy the day before you start HEMLIBRA
- You may continue taking your prophylactic factor VIII for the first week of HEMLIBRA

HEMLIBRA may interfere with laboratory tests that measure how well your blood is clotting and create an inaccurate result. Speak with your healthcare provider about how this may affect your care.

These are not all of the possible side effects of HEMLIBRA. Speak to your healthcare provider for medical advice about side effects.

Side effects may be reported to the FDA at (800) FDA-1088 or www.fda.gov/medwatch. You may also report side effects to Genentech at (888) 835-2555.

Please see Important Safety Information, including **Serious Side Effects**, throughout this brochure, as well as the accompanying HEMLIBRA full Prescribing Information and Medication Guide.





IMPORTANT INFORMATION BEFORE YOU GET STARTED

Before injecting HEMLIBRA, it's important that you read, understand, and follow the Instructions for Use. Your healthcare provider should show you, or your caregiver, how to prepare, measure, and inject HEMLIBRA properly before you use it for the first time. Ask your healthcare provider if you have any questions.

IMPORTANT INFORMATION

- **Do not** inject yourself or someone else unless you have been shown how to by your healthcare provider
- Check the expiration date on the box and vial label. **Do not** use if the expiration date has passed
- Make sure the name HEMLIBRA appears on the box and vial label
- **Only use the vial one time.** After you inject your dose, dispose of (throw away) any unused HEMLIBRA left in the vial. Do not save unused HEMLIBRA in the vial for later use
- Before opening the vial, read the vial label to make sure you have the medicine strength(s) needed to give the dose prescribed by your healthcare provider
- **Only use the syringes, transfer needles, and injection needles that your healthcare provider prescribes**
- Your healthcare provider will determine your dose in milliliters (mL) that you will need to give based on your body weight
- **Only use the syringes, transfer needles, and injection needles one time. Dispose of (throw away) any used syringes and needles in a sharps disposal container**
- HEMLIBRA comes in 4 strengths. Depending on your dose, you may need to use more than one vial to give your total prescribed dose. **Do not combine HEMLIBRA vials of different concentrations in one injection** to give the prescribed dose
- If your prescribed dose is more than 2 mL, you will need to give more than one injection of HEMLIBRA

Please see Important Safety Information, including **Serious Side Effects**, throughout this brochure, as well as the accompanying HEMLIBRA full Prescribing Information and Medication Guide.


HEMLIBRA
emicizumab-kxwh | 150
injection for subcutaneous use | mg/mL

Some of the supplies necessary for your injection may be familiar, while others may be new to you.

ITEMS INCLUDED IN THE HEMLIBRA BOX



Available HEMLIBRA vials

ITEMS NOT INCLUDED IN THE HEMLIBRA BOX



ALCOHOL WIPES



GAUZE



25- TO 27-GAUGE INJECTION
NEEDLE WITH SAFETY SHIELD

Used to inject HEMLIBRA;
do not use the injection needle
to withdraw HEMLIBRA from vial.



18-GAUGE TRANSFER NEEDLE
WITH 5 MICRON FILTER

If you need to use more than one
vial to inject your prescribed dose,
you must use a new transfer needle
for each vial. Do not use the transfer
needle to inject HEMLIBRA.



SYRINGE

For injection amount up to 1 mL,
use a 1 mL syringe. For injection
amount between 1 mL and 2 mL,
use a 2 mL or 3 mL syringe.



COTTON BALLS



SHARPS DISPOSAL
CONTAINER

IS AN INJECTION MAT
NEEDED?

You may want to use the injection mat, which came in your HEMLIBRA Welcome Kit. While an injection mat is not required, it may be helpful to organize supplies for your HEMLIBRA injection. If you didn't receive an injection mat and would like to order one, or if you misplaced yours and would like a new one, call us at **1-866-HEMLIBRA** (1-866-436-5427).

You may need two of some items, depending on the dose that your healthcare provider prescribes. If your healthcare provider prescribes a dose that is more than 2 mL, you'll need to give more than one injection. Only use the syringes, transfer needles, and injection needles that your healthcare provider prescribes. You may get these supplies from your specialty pharmacy or your healthcare provider's office.



PREPARING FOR
YOUR INJECTION

STORING HEMLIBRA



You should keep your HEMLIBRA in the refrigerator at 36°F to 46°F (2°C to 8°C). Before giving an injection, unopened vials of HEMLIBRA may be stored out of the refrigerator and then returned to the refrigerator. Be sure not to store HEMLIBRA out of the refrigerator for more than a total of 7 days or at a temperature greater than 86°F (30°C).

It's important to keep your HEMLIBRA in its original package to protect it from light until you are ready to use it. Be sure not to freeze or shake HEMLIBRA. Keep HEMLIBRA and all medicines out of the reach of children.

INSPECTING HEMLIBRA



Remember to inspect all supplies before you start, including the box and vial label. If they are damaged, have been dropped, or are expired, don't use them; get another set of supplies. Once ready, place the supplies on a clean, well-lit, flat work surface.

Double check to make sure the name HEMLIBRA appears on the box and vial label. Also, check your medication to be sure it is the right strength, isn't damaged, and hasn't expired.

You'll also want to check the appearance of your medication. HEMLIBRA is a colorless to slightly yellow liquid. Don't use it if it is damaged, looks cloudy, hazy, or colored, is missing the cap covering the stopper, or has particles.


SELECTING YOUR INJECTION SITE



Before you start, you will need to decide where to inject. You should use a different injection site each time you give an injection, at least 1 inch away from the area you used for your previous injection.

You can use your:

- Thigh (front or middle)
- Stomach area (abdomen), except for the 2 inches around the belly button
- Outer area of the upper arm (only if a trained caregiver is giving the injection)



Do not inject into areas that could be irritated by a belt or waistband.

Do not inject into moles, scars, bruises, or areas where the skin is tender, red, hard, or broken.

Once you select an injection site, clean the chosen area using an alcohol wipe. Let the skin dry for about 10 seconds. Do not touch, fan, or blow on the cleaned area before your injection.

KEEPING TRACK OF YOUR INJECTION



You may find it useful to keep track of your injections and your injection sites, in a log that you can share with your healthcare provider as needed. You may wish to use the log provided in the HEMLIBRA Welcome Kit or continue using the system that works well for you today.

PREPARING YOUR MEDICINE



Before injecting your medicine, make sure to take the vial(s) out of the refrigerator 15 minutes before use, and allow it to reach room temperature on a clean, flat surface away from direct sunlight. Do not try to warm the vial by any other way. Make sure to wash your hands with soap and water.



Please see Important Safety Information, including **Serious Side Effects**, throughout this brochure, as well as the accompanying HEMLIBRA full Prescribing Information and Medication Guide.


TRANSFERRING HEMLIBRA

FROM VIAL TO SYRINGE

STEP 1. REMOVE VIAL CAP AND CLEAN TOP




- Take the cap off the vial(s).




- Clean the top of the vial(s) stopper with an alcohol wipe.
- Dispose of (throw away) the vial cap(s) into the sharps disposal container.

STEP 2. ATTACH TRANSFER NEEDLE WITH FILTER TO SYRINGE




- Push and twist the transfer needle with filter clockwise onto the syringe until it is fully attached.




- Slowly pull back on the plunger and draw air into the syringe that is the same amount as your prescribed dose.

STEP 3. UNCAP TRANSFER NEEDLE




- Hold the syringe by the barrel with the transfer needle pointing up.
- Carefully pull the transfer needle cap straight off and away from your body. **Do not throw the cap away. Place the transfer needle cap down on a clean, flat surface.** You will need to recap the transfer needle after transferring the medicine.
- **Do not touch** the needle tip or place it on a surface after the needle cap has been removed.


STEP 4. INJECT AIR INTO VIAL



- Keep the vial on the flat working surface and insert the transfer needle and syringe straight down into the **center** of the vial stopper.



- Keep the needle in the vial and turn the vial upside down.




- With the needle pointing upwards, push on the plunger to inject the air from the syringe **above the medicine**.
- Keep your finger pressed down on the syringe plunger.
- **Do not** inject air into the medicine.

TRANSFERRING HEMLIBRA

FROM VIAL TO SYRINGE



STEP 5. TRANSFER HEMLIBRA TO SYRINGE



- Slide the tip of the needle down so that it is **within the medicine**.
- Slowly pull back the plunger to fill the syringe with more than the amount of HEMLIBRA needed for your prescribed dose.
- Be careful not to pull the plunger out of the syringe.

Important: If your prescribed dose is more than the amount of HEMLIBRA in the vial, **withdraw all HEMLIBRA** and go to the **"Combining Vials"** section on page 17.


STEP 6. REMOVE AIR BUBBLES



- Keep the needle in the vial and check the syringe for larger air bubbles. Too large an air bubble can reduce the dose you receive.
- **Remove the larger air bubbles** by gently tapping the syringe barrel with your fingers until the air bubbles rise to the top of the syringe. Move the tip of the needle **above the medicine** and slowly push the plunger up to push the air bubbles out of the syringe.
- If the amount of HEMLIBRA in the syringe is now at or below your prescribed dose, move the tip of the needle to **within the medicine** and slowly **pull** back the plunger until you have **more** than the amount of HEMLIBRA needed for your **prescribed dose**.
- Be careful not to pull the plunger out of the syringe.
- Repeat the steps above until you have removed the larger air bubbles.


Note: Ensure you have enough HEMLIBRA in the syringe to complete your dose before moving on to the next step. If you cannot remove all of HEMLIBRA, turn the vial upright to reach the remaining amount.

STEP 7. RECAP TRANSFER NEEDLE




- Remove the syringe and transfer needle from the vial.
- **Using one hand, slide** the transfer needle into the cap and **scoop upwards** to cover the needle.
- Once the needle is covered, push the transfer needle cap towards the syringe to fully attach it with **one hand** to prevent accidentally sticking yourself with the needle.

STEP 8. CLEAN INJECTION SITE



- Select and **clean** your injection site with an alcohol wipe.
- Let the skin dry for about 10 seconds. Do not touch, fan, or blow on the cleaned area before your injection.

STEP 9. REMOVE TRANSFER NEEDLE



- Remove the transfer needle from the syringe by twisting counter-clockwise and gently pulling.
- Dispose of (throw away) the used transfer needle into a sharps disposal container.
- **Note:** Please proceed to the next page for further instructions on how to inject.


Do not use the transfer needle to inject HEMLIBRA, as this may cause harm such as pain and bleeding.

INJECTING
HEMLIBRA

Now that you've prepared your syringe and cleaned your injection site, it's time to inject HEMLIBRA.


Please read the instructions fully and follow them step-by-step. Your healthcare provider will help you through the process to make sure you feel comfortable before you inject for the first time. You can also refer to the Instructions for Use, included in the pocket of this brochure, for any additional questions.

STEP 10. ATTACH INJECTION NEEDLE TO SYRINGE




- Push and twist the injection needle clockwise onto the syringe until it is fully attached.

STEP 11. MOVE SAFETY SHIELD



- Move the safety shield away from the needle and **towards** the syringe barrel.

STEP 12. UNCAP INJECTION NEEDLE




- Carefully pull the injection needle cap away from the syringe.
- Dispose of (throw away) the cap into a sharps disposal container.
- **Do not touch** the needle tip or allow it to touch any surface.
- After the injection needle cap has been removed, HEMLIBRA in the syringe must be injected right away.

Note: HEMLIBRA must not be stored in the syringe. HEMLIBRA in the syringe must be injected under the skin (subcutaneous injection) immediately.


Please see Important Safety Information, including **Serious Side Effects**, throughout this brochure, as well as the accompanying HEMLIBRA full Prescribing Information and Medication Guide.

STEP 13. ADJUST PLUNGER TO PRESCRIBED DOSE



- Slowly push the plunger to your prescribed dose.
- Ensure the top rim of the plunger is in line with the mark on the syringe for your prescribed dose.


STEP 14. SUBCUTANEOUS (UNDER THE SKIN) INJECTION



- Pinch the selected injection site and fully insert the needle at a **45° to 90°** angle with a quick, firm action. **Do not** hold or push on the plunger while inserting the needle.
- Hold the position of the syringe and let go of the pinched injection site.


Please see Important Safety Information, including **Serious Side Effects**, throughout this brochure, as well as the accompanying HEMLIBRA full Prescribing Information and Medication Guide.

STEP 15. INJECT HEMLIBRA




- Slowly inject all of HEMLIBRA by gently pushing the plunger all the way down.
- Remove the needle and syringe from the injection site at the same angle as inserted.

STEP 16. COVER NEEDLE WITH SAFETY SHIELD



- Move the safety shield forward 90°, away from the syringe barrel.
- Holding the syringe with one hand, **press the safety shield down** against a flat surface with a firm, quick motion until you hear a "click."




- If you do not hear a click, look to see that the needle is fully covered by the safety shield.
- Keep your fingers behind the safety shield and away from the needle at all times.
- **Do not** remove the injection needle from the syringe.

You may need to use more than one vial to inject your total prescribed dose.

If you do, follow the steps below after you have drawn up HEMLIBRA from the first vial. **Do not combine HEMLIBRA vials of different concentrations in one injection to give the prescribed dose.**


If you have any questions, please ask your healthcare provider.

STEP A. RECAP TRANSFER NEEDLE




- Remove the syringe and transfer needle from the first vial.
- **Using one hand**, slide the transfer needle into the cap and **scoop upwards** to cover the needle.
- Once the needle is covered, push the transfer needle cap toward the syringe to fully attach it with **one hand** to prevent accidentally sticking yourself with the needle.

STEP B. REMOVE TRANSFER NEEDLE




- Remove the transfer needle from the syringe by twisting counter-clockwise and gently pulling.
- Dispose of (throw away) the used transfer needle into a sharps disposal container.

STEP C. ATTACH A NEW TRANSFER NEEDLE WITH FILTER TO SYRINGE




- Push and twist a **new** transfer needle with filter clockwise onto the syringe until it is fully attached.
- Slowly pull back the plunger and draw some air into the syringe.
- **Note: You must use a new transfer needle each time you withdraw HEMLIBRA from a new vial.**

STEP D. UNCAP TRANSFER NEEDLE




- Hold the syringe by the barrel with the transfer needle cap pointing up.
- Carefully pull the transfer needle cap straight off and away from your body. **Do not throw the cap away.** You will need to recap the transfer needle after drawing up the medicine.
- **Do not touch** the needle tip.


STEP E. INJECT AIR INTO VIAL



- With the new vial on the flat working surface, insert the new transfer needle and syringe, straight down into the **center** of the vial stopper.




- Keep the transfer needle in the vial and turn the vial upside down.



- With the needle pointing upwards, inject the air from the syringe **above the medicine**.
- Keep your finger pressed down on the syringe plunger.
- **Do not** inject air into the medicine.

STEP F. TRANSFER HEMLIBRA TO SYRINGE



- Slide the tip of the needle down so that it is **within the medicine**.
- Slowly pull back the plunger to fill the syringe barrel more than the amount of HEMLIBRA needed for your prescribed dose.
- Be careful not to pull the plunger out of the syringe.
- **Do not** use the transfer needle to inject HEMLIBRA as this may cause harm such as pain and bleeding.
- **Note:** Ensure you have enough HEMLIBRA in the syringe to complete your dose before moving on to the next steps. If you cannot remove all of HEMLIBRA, turn the vial upright to reach the remaining amount.


HOW DO I COMBINE THE VIALS?

Repeat steps A to F with each additional vial until you have more than the amount of HEMLIBRA needed for your prescribed dose. Once completed, keep the transfer needle inserted in the vial and return to Transfer Needle Preparation Step 6 **"Remove Air Bubbles"** on page 12. Continue with the remaining steps.

DISPOSING OF YOUR SUPPLIES

Once you've finished injecting HEMLIBRA, dispose of any used vial(s), needles, injection needle caps, and used syringes in a sharps disposal container. **Do not** dispose of (throw away) any loose caps, vials, needles, or syringes in your household trash.

If you do not have an FDA-cleared sharps disposal container, you may use a household container that is made of heavy-duty plastic; can be closed with a tight-fitting, puncture resistant lid, without sharps being able to come out; remains upright and stable during use; is leak-resistant; and is properly labeled to warn of hazardous waste inside the container.



- **Only use the vial one time.** Throw away any unused medicine in the vial. Do not save unused medicine for later use.
- **Do not** try to remove the used injection needle from the used syringe.
- **Do not recap** the injection needle with the cap.
- Do not dispose of (throw away) any used sharps disposal container in your household trash unless your community guidelines permit this. Do not recycle your used sharps disposal container.
- When your sharps disposal container is almost full, you will need to follow your community guidelines for the right way to dispose of (throw away) your sharps disposal container. For more information about safe sharps disposal and for specific information about sharps disposal in the state that you live in, go to the FDA's website at <http://www.fda.gov/safesharpsdisposal>.
- **Important: Always keep the sharps disposal container out of reach of children.**

WHAT DO I DO AFTER INJECTION?

Do not rub the injection site after your injection. You may see a few drops of blood. If you do, just press a cotton ball or gauze on it to stop the bleeding. If you experience any bruising, you can also use an ice pack. If bleeding does not stop, contact your healthcare provider.

Please see Important Safety Information, including **Serious Side Effects**, throughout this brochure, as well as the accompanying HEMLIBRA full Prescribing Information and Medication Guide.

SUMMARY OF
INJECTION TIPS

Below is a list of tips to remember before and after injecting HEMLIBRA. For a full list of instructions, please refer to the HEMLIBRA Instructions for Use.

- **Leave your HEMLIBRA out for at least 15 minutes** to bring it to room temperature before injecting. (For example, you may take HEMLIBRA out the morning of your injection to let it get to room temperature as you get ready for the day or take HEMLIBRA out before you go to bed at night if you plan to take your injection first thing in the morning)
- If you are nervous about your first injection, it may be helpful to **use some relaxation techniques or video chat with a friend for support**
- Make sure to **rotate injection sites** each time you treat. They should be at least 1 inch from the previous injection site
- To avoid bubbles or foaming, **slowly** pull the plunger back when filling the syringe
- After using an alcohol swab to clean the injection site, **let the skin dry for at least 10 seconds** before injecting
- **Pinch the skin** to avoid muscle when injecting
- **The medication should be injected slowly** after the needle has penetrated the skin
- After the injection, **do not rub the injection site**
- If there is bruising, **an ice pack can be applied** with gentle pressure
- Always **talk to your doctor** if you have questions about how to administer HEMLIBRA, including any other injection tips

Tips to help you stay on track

It is important to take HEMLIBRA exactly as prescribed by your healthcare provider. Here are some tips to help you stick to your prescribed dosing routine.



Choose your injection day and time; they should remain the same for each injection.




Set reminders on your phone or computer calendar or ask your healthcare provider about helpful apps.



Use a wall or desk calendar to mark your injection day; you may want to use the calendar reminder stickers provided in the injection log in the HEMLIBRA Welcome Kit.

Consider injection tracking tools

You may find it useful to keep track of your injections in a log that you can share with your healthcare provider as needed.



You may wish to use the log provided in the HEMLIBRA Welcome Kit, or continue using the system that works well for you today. As you record your injection, remember to:

- Write down the date, time, and vial lot information in case you need it for your healthcare provider at some point in the future
- Write down the spot you used for injection so that you don't forget
- Track if your supplies are running low. If they are, contact your healthcare provider to order more so that you always have what you need to inject HEMLIBRA

INJECTION SUPPORT

RESOURCES

If you need a refresher on how to inject HEMLIBRA, you can:

- Contact your healthcare provider with any questions
- Review the Instructions for Use that came with your HEMLIBRA
- Visit TakingHEMLIBRA.com to watch injection videos
- Contact your local Genentech Account and Community Manager (ACM), who can provide education following training by your healthcare provider. Do not attempt to inject yourself or another person unless you have been taught how to do so by a healthcare provider. ACMs are Genentech employees and do not provide medical advice. Always talk to your doctor about treatment options

• • • • **CONNECTS**

PATIENT RESOURCE CENTER

CALL (866) HEMLIBRA OR (866) 436-5427.

Our comprehensive resource hub, HEMLIBRA Connects, helps get you the information you need about HEMLIBRA. We can help you:

- Be connected to your local Account and Community Manager
- Learn about our HEMLIBRA Co-Pay Program that may help you with the out-of-pocket costs of HEMLIBRA, if you are eligible
- Understand your health insurance coverage for HEMLIBRA

HEMLIBRA Access Solutions® may be able to help you understand how to get the HEMLIBRA you need, even if you can't pay for it. Find out more in the Access Solutions brochure in the HEMLIBRA Welcome Kit or by visiting HEMLIBRA.com/ACCESS

We're here to help, in English and Spanish, 9:00 AM to 8:00 PM ET (6:00 AM to 5:00 PM PT), Monday through Friday.

Sign up at HEMLIBRA.com to receive news updates and treatment tips.

The information contained in this brochure does not provide medical advice or take the place of speaking with your healthcare provider about your medical condition or treatment.

Please see Important Safety Information, including Serious Side Effects, throughout this brochure, as well as the accompanying HEMLIBRA full Prescribing Information and Medication Guide.



HEMLIBRA® is a registered trademark of Chugai Pharmaceutical Co., Ltd., Tokyo, Japan.
The HEMLIBRA logo is a registered trademark of Chugai Pharmaceutical Co., Ltd., Tokyo, Japan.
The Genentech logo is a registered trademark of Genentech, Inc.
All other trademarks are the property of their respective owners.
©2023 Genentech USA, Inc. All rights reserved. M-US-00002705(v6.0) 07/23

