

IS HEMLIBRA®

RIGHT FOR YOU?



A GUIDE TO HELP YOU TALK TO
YOUR OR YOUR CHILD'S DOCTOR.

PROPHYLAXIS FOR ALL PEOPLE WITH HEMOPHILIA A

You know that managing hemophilia A can be complicated. HEMLIBRA is the only subcutaneous injection and prophylactic treatment for adults and children, ages newborn and older, with hemophilia A with or without factor VIII inhibitors. These questions can help you and your doctor decide if HEMLIBRA may be right for you or your child.

In the past year, how many bleeds did you have with your current treatment?

0 – 1 – 2 – 3 – 4 – 5 – 6 – 7 – 8 – 9 – 10 – 11+

How many of those did you treat?

How many times a week do you take your current treatment?

1 – 2 – 3 – 4 – 5 – 6 – 7

How comfortable are you traveling, playing sports, or participating in other physical activities while on your current treatment?

1 ————— 2 ————— 3 ————— 4 ————— 5
Very uncomfortable Neutral Extremely comfortable

In the last year, have you gone to the ER because of a bleed? If so, how many times?

☐ Yes ☐ No # of times

How well are you adhering to your current treatment plan?

1 ————— 2 ————— 3 ————— 4 ————— 5
Not at all Somewhat Completely

What are your goals with your hemophilia A treatment?

What is HEMLIBRA?

HEMLIBRA is a prescription medicine used for routine prophylaxis to prevent or reduce the frequency of bleeding episodes in adults and children, ages newborn and older, with hemophilia A with or without factor VIII inhibitors.

What is the most important information I should know about HEMLIBRA?

HEMLIBRA increases the potential for your blood to clot. People who use activated prothrombin complex concentrate (aPCC; Feiba®) to treat breakthrough bleeds while taking HEMLIBRA may be at risk of serious side effects related to blood clots.

These serious side effects include:

- **Thrombotic microangiopathy (TMA)**, a condition involving blood clots and injury to small blood vessels that may cause harm to your kidneys, brain, and other organs
- **Blood clots (thrombotic events)**, which may form in blood vessels in your arm, leg, lung, or head

Carefully follow your healthcare provider's instructions regarding when to use an on-demand bypassing agent or factor VIII, and the dose and schedule to use for breakthrough bleed treatment. If aPCC (Feiba®) is needed, talk to your healthcare provider in case you feel you need more than 100 U/kg of aPCC (Feiba®) total.

Your body may make antibodies against HEMLIBRA, which may stop HEMLIBRA from working properly. Contact your healthcare provider immediately if you notice that HEMLIBRA has stopped working for you (eg, increase in bleeds).

Please see Important Safety Information, including **Serious Side Effects**, throughout this piece, as well as the HEMLIBRA full [Prescribing Information](#) and [Medication Guide](#).

YOUR HEMLIBRA®

CONVERSATION STARTS NOW

Appointment Date/Time

/ /

: AM
PM

Choosing the right treatment is an important decision that involves careful consideration. When you're ready to talk to your doctor, these questions can help get you started.

How is HEMLIBRA different from my current treatment?

What is half-life, and why does that matter?

How often do I need to take HEMLIBRA?

Do I need to take factor VIII or bypassing agents while I'm taking HEMLIBRA?

What are the side effects of HEMLIBRA?

Can I get a bleed while taking HEMLIBRA?

SCHEDULE YOUR DOCTOR'S APPOINTMENT TODAY. TOGETHER WITH THIS GUIDE, YOU CAN DECIDE IF HEMLIBRA IS RIGHT FOR YOU.

Important Safety Information (continued)

What are the other possible side effects of HEMLIBRA?

The most common side effects of HEMLIBRA include: injection site reactions (redness, tenderness, warmth, or itching at the site of injection), headache, and joint pain. These are not all of the possible side effects of HEMLIBRA. You can speak with your healthcare provider for more information.

What else should I know about HEMLIBRA?

See the detailed "Instructions for Use" that comes with your HEMLIBRA for information on how to prepare and inject a dose of HEMLIBRA, and how to properly throw away (dispose of) used needles and syringes.

- Stop taking your prophylactic bypassing therapy the day before you start HEMLIBRA
- You may continue taking your prophylactic factor VIII for the first week of HEMLIBRA

HEMLIBRA may interfere with laboratory tests that measure how well your blood is clotting and create an inaccurate result. Speak with your healthcare provider about how this may affect your care.

These are not all of the possible side effects of HEMLIBRA. Speak to your healthcare provider for medical advice about side effects.

Side effects may be reported to the FDA at (800) FDA-1088 or www.fda.gov/medwatch. You may also report side effects to Genentech at (888) 835-2555.

Please see Important Safety Information, including **Serious Side Effects**, throughout this piece, as well as the HEMLIBRA full [Prescribing Information](#) and [Medication Guide](#).

Genentech

A Member of the Roche Group

HEMLIBRA® is a registered trademark of Chugai Pharmaceutical Co., Ltd., Tokyo, Japan.

The HEMLIBRA logo is a trademark of Chugai Pharmaceutical Co., Ltd., Tokyo, Japan.

The Genentech logo is a registered trademark of Genentech, Inc.

© 2023 Genentech USA, Inc. All rights reserved. For U.S. residents only. M-US-00001641(v3.0) 01/23


HEMLIBRA®
emicizumab-kxwh | 150 mg/mL
injection for subcutaneous use